



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Bo
Search	PubMed	▼	for	schubert, w.			Go	Clear
Limits		Preview/Index		History		Clipboard		Details

Display	Summary	▼	Show:	20	▼	Sort	▼	Send to	Text	▼
Items 1-2 of 2										One page

Entrez PubMed

- ☐ 1: [Kamlowski A, van der Est A, Fromme P, Krauss N, Schubert WD, Klukas O, Stehlik D.](#) [Related Articles, Links](#)



The structural organization of the PsaC protein in Photosystem I from single crystal EPR and X-ray crystallographic studies.
Biochim Biophys Acta. 1997 Apr 11;1319(2-3):199-213.
PMID: 9131044 [PubMed - indexed for MEDLINE]

PubMed Services

- ☐ 2: [Schubert W.](#) [Related Articles, Links](#)



["Transfusion hepatitis following exchange transfusion" published in issue 22-1972 of this journal. Reply to the article of H. Scholz, W. Dorffel and Inge Mitschke]
Dtsch Gesundheitsw. 1972 Dec 7;27(49):2347. German. No abstract available.
PMID: 4569684 [PubMed - indexed for MEDLINE]

Related Resources

[Write to the Help Desk](#)



[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

Aug 6 2003 12:56:11

0



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Bo	
Search	PubMed	▼	for					Go	Clear
Limits		Preview/Index		History		Clipboard		Details	

Display	Summary	▼	Show	20	▼	Sort	▼	Send to	Text	▼
Items 1-4 of 4										One page

Entrez PubMed

☐ 1: [Jain KK.](#)[Related Articles, Links](#)

Applications of proteomics in oncology.
Pharmacogenomics. 2000 Nov;1(4):385-93. Review.
PMID: 11257924 [PubMed - indexed for MEDLINE]

☐ 2: [Marshall T, Williams KM.](#)[Related Articles, Links](#)

Proteomics and its impact upon biomedical science.
Br J Biomed Sci. 2002;59(1):47-64. Review.
PMID: 12000188 [PubMed - indexed for MEDLINE]

PubMed Services

☐ 3: [Bock JR, Gough DA.](#)[Related Articles, Links](#)

Predicting protein--protein interactions from primary structure.
Bioinformatics. 2001 May;17(5):455-60.
PMID: 11331240 [PubMed - indexed for MEDLINE]

Related Resources


☐ 4: [Song EJ, Lee KJ.](#)[Related Articles, Links](#)

[Identification of proteome molecules by proteomics using
two-dimensional gel electrophoresis and MALDI-TOF MS]
Exp Mol Med. 2001 Apr 21;33(1 Suppl):5-18. Korean.
PMID: 11708325 [PubMed - indexed for MEDLINE]

[Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act | Disclaimer](#)

Aug 6 2003 12:56:11

0



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Bo	
Search	PubMed	▼	for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details			
Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼

☐ 1: Adv Biochem Eng Biotechnol. 2003;83:189-209.

[Related Articles, Links](#)

Entrez PubMed

Topological proteomics, toponomics, MELK-technology.

Schubert W.

PubMed Services

MelTec Ltd., ZENIT-Building, Leipziger Strasse 44, 39120 Magdeburg, Germany. info@meltec.de

Related Resources



MELK is an ultrasensitive topological proteomics technology analysing proteins on the single cell level (Multi-Epitope-Ligand-'Kartographie'). It can trace out large scale protein patterns with subcellular resolution, mapping the topological position of many proteins simultaneously in a cell. Thereby, it addresses higher level order in a proteome, referred to as the toponome, coding cell functions by topologically and timely determined webs of interacting proteins. The resulting cellular protein maps provide new structures in the proteome: single combinatorial protein patterns (s-CPP), and combinatorial protein pattern motifs (CPP-motifs), bound to superior units. They are images of functional protein networks, which are specific signatures of tissues, cell types, cell states and diseases. The technology unravels hierarchies of proteins related to particular cell functions or dysfunctions, thus identifying and prioritising key proteins within cell and tissue protein networks. Interlocking MELK with the drug screening machinery provides new clues related to the selection of target proteins, and functionally relevant hits and drug leads. The present chapter summarizes the steps that have contributed to the establishment of the technology.

PMID: 12934931 [PubMed - in process]

Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	----------	---	------	---	---------	------	---

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Aug 6 2003 12:56:11



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Bo	
Search	PubMed	▼	for					Go	Clear
Limits		Preview/Index		History		Clipboard		Details	

Display	Abstract	▼	Show:	20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	-------	----	---	------	---	---------	------	---

☐ 1: Dtsch Gesundheitsw. 1972 Dec 7;27(49):2347.

[Related Articles, Links](#)

Entrez PubMed

**["Transfusion hepatitis following exchange transfusion"
published in issue 22-1972 of this journal. Reply to the article
of H. Scholz, W. Dorffel and Inge Mitschke]**

PubMed Services

[Article in German]

Schubert W.

Publication Types:

- Biography
- Historical Article

Personal Name as Subject:

- Scholz H
- Dorffel W
- Mitschke I

Related Resources

PMID: 4569684 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Show:	20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	-------	----	---	------	---	---------	------	---

[Write to the Help Desk](#)



[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

Aug 6 2003 12:56:11

0



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Bo
Search	PubMed	for					Go	Clear
Limits		Preview/Index		History		Clipboard		Details
Display	Abstract		Show: 20		Sort		Send to	Text

Entrez PubMed

☐ 1: Biochim Biophys Acta. 1997 Apr
11;1319(2-3):199-213.

[Related Articles, Links](#)

The structural organization of the PsaC protein in Photosystem I from single crystal EPR and X-ray crystallographic studies.

PubMed Services

Kamlowski A, van der Est A, Fromme P, Krauss N, Schubert WD, Klukas O, Stehlik D.

Institut fur Experimentalphysik, Freie Universitat Berlin, Germany.



Related Resources

In Photosystem I (PS I) the terminal electron acceptors, FA and FB, are iron-sulfur (4Fe-4S) centers, which are bound to the stromal subunit PsaC. The orientation of PsaC is determined relative to the whole PS I complex (see Schubert, W.-D. et al. (1995) in From Light to Biosphere (Mathis, P. ed.), Vol. II, pp. 3-10, Kluwer) from which a molecular model for the structure of PsaC within PS I is derived. Two strategies are followed: (i) PS I single crystal EPR data on the orientation of the g tensors of both FA- and FB- relative to each other and relative to the crystal axes (see preceding paper) are used in conjunction with the central structural part of the bacterial 2 [Fe4S4] ferredoxins, the cysteine binding motifs of which are known to be homologous to those of PsaC; (ii) the same core structure is fitted into the intermediate resolution electron density map of PS I. The PsaC orientation obtained both ways agree well. The local twofold symmetry axis inherent to the ferredoxin model leaves a twofold ambiguity in the structural conclusion. Deviations from this C2-symmetry in the amino acid sequence of PsaC are analyzed with respect to observable properties which would resolve the remaining structural ambiguity. Arguments both for and against FA being the distal iron-sulfur center (to FX) are discussed.

PMID: 9131044 [PubMed - indexed for MEDLINE]

Display	Abstract		Show: 20		Sort		Send to	Text
---------	----------	--	----------	--	------	--	---------	------

[Write to the Help Desk](#)



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Bo
Search PubMed	for						Go	Clear
Limits		Preview/Index		History		Clipboard		Details
Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text

☐ 1: Exp Mol Med. 2001 Apr 21;33(1 Suppl):5-18.

[Related Articles, Links](#)

Entrez PubMed

FREE full text article at
www.e-9mm.org

[Identification of proteome molecules by proteomics using two-dimensional gel electrophoresis and MALDI-TOF MS]

PubMed Services

[Article in Korean]


Song EJ, Lee KJ.

College of Pharmacy and Division of Molecular Life Sciences Ewha Womans University, Seoul, Korea.

Related Resources

Genomic technologies have enabled rapid accumulation of information from complex biological systems over the last two decades. The complete DNA sequence is now known for many organisms and the informational database obtained from genome sequencing projects has provided the base for the specification of proteome - the protein complement of genome. Genomic functions can be inferred from the analysis of gene structure and gene expression profiles because proteins are the functional molecules of an organism. Integrated technologies including protein separation, identification, characterization and information manage system are essential to analyze the proteins in complex cellular matrix. This study is focusing on the strategies of proteome analysis using sample preparation, 2-dimensional gel electrophoresis, processing of protein spots and identification of proteins, protein-protein interaction and posttranslational modification using MALDI-TOF-MS. 2-D gel electrophoresis is currently the most powerful protein separation technique and MALDI-TOF MS is powerful identification technique for protein and peptides as a sensitive, rapid, and high resolution analytical method. The developed integrated proteome technologies are very useful to understand the biological phenomena at molecular level by identifying the new molecules and their modifications in various cellular processes, and can be applied for biotechnology including medical science.

PMID: 11708325 [PubMed - indexed for MEDLINE]

nature**nature** 26 April 1990 [home](#) [search](#) [help](#)**letters to nature***Nature* **344**, 868 - 870 (1990); doi:10.1038/344868a0**Activin is a nerve cell survival molecule**

D. Schubert, H. Kimura, M. LaCorbiere, J. Vaughan, D. Karr & W. H. Fischer

contents**pdf**

THE structures of five neurotrophic molecules have so far been published. Nerve growth factor¹, fibroblast growth factor^{2, 3} and purpurin⁴, have been identified as nerve-cell survival molecules. More recently, brain-derived neurotrophic factor (BDNF) and ciliary neurotrophic factor have been cloned and sequenced^{5, 6}. As all these proteins stimulate the survival of ciliary or sensory neurons, a new cell survival assay is required if novel neurotrophic molecules are to be discovered. P19 teratoma cells differentiate to nerve-like cells in the presence of 5×10^{-7} M retinoic acid (RA)^{7, 8}. But when P19 cells are plated in N² synthetic medium⁹ without being exposed to RA, they die within 48 h. In an attempt to identify a molecule(s) that can substitute for RA in promoting P19 survival, we assayed serum-free growth-conditioned media for their ability to promote P19 survival. One cell line from the rat eye secreted a molecule that promoted the survival of P19 cells and some types of nerve cell. We identified this molecule as activin, better known for its role in hormone secretion.



Nature © Macmillan Publishers Ltd 1990 Registered No. 785998

England.